



Psychophysiological Profiles of Sleep Deprivation and Stress during Marine Corps Training

Chris Berka¹, Gene Davis¹, Robin Johnson¹, Daniel J. Levendowski¹, Melissa Whitmoyer¹, Robin Fatch¹, Wayne Ensign², Matthew A. Yanagi², and Richard Olmstead³

**(1) Advanced Brain Monitoring, Inc., Carlsbad, CA,
(2) SPAWAR, San Diego, CA,
(3) VA Greater Los Angeles Healthcare System**

Research supported by the Defense Advanced Research Projects Agency (DARPA) program “Preventing Sleep Deprivation”, and by the Office of Naval Research. DARPA has approved this material for public release, distribution unlimited.

Technology development also funded by NIH.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 2007		2. REPORT TYPE		3. DATES COVERED 00-00-2007 to 00-00-2007	
4. TITLE AND SUBTITLE Psychophysiological Profits of Sleep Deprivation and Stress during Marine Corps Training				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Advanced Brain Monitoring Inc,2237 Faraday Ave Ste 100,Carlsbad,CA,92008				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES in Sleep30:A132					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 16	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			



Introduction: Study Goals & Rationale

- Assess fatigue and stress and their effects on situation awareness and decision-making during 28-day USMC Mojave Viper training
- Evaluate utility of EEG, EKG, actigraphy in predicting performance deficits in an operational setting

Value to USMC

Better awareness of team capabilities

Improve safety and mission management

Fatigue management



Challenges of the operational environment

Mojave Viper training at 29 Palms

- 28-day training event covering all USMC operations in desert & urban environments
- Final training prior to deployment in Iraq or Afghanistan

Main objectives

1. Demonstrate that marines experience fatigue during training using objective measures.
2. Assess effects of fatigue and/or stress on performance during training.



Methods: participants & measures

- N=17 platoon, squad & fire team leaders
- Continuous actigraphy throughout testing
- Baseline and repeated sessions (1/wk)
 - 30-minute EEG, EKG, 3-Choice Vigilance
 - Saliva cortisol levels
 - Subjective measures
 - POMS
 - Brief Fatigue Inventory
 - Perceived Stress Scale
 - Subjective Sleepiness Scale




Mobile Wireless EEG System




No-prep sensors
Battery-powered
Automate artifact
detect / decon

- EEG sensors at Fz, Cz, POz referenced to linked mastoids, bipolar Fz-PO and Cz-PO
- EKG sensors on collarbone & lower rib
- System proved durable in field environment
 - ✓ Marines quickly learned to self-apply
 - ✓ No issues with comfort or usability








Results: Actigraphy- Sleep Debt



- Sleep debt calculated by subtracting recommended sleep period (8 h/ 480 min) from actual sleep minutes.
- By night 21, the marines average 58 hours of accumulated sleep debt




Results: 3-Choice Vigilance Test






Results: EEG-based Fatigue indications

Linear Drowsiness Scale




Anchoring LDS





Results: Heart rate/variability



HR/HRV measures

Calculated on 5 min. windows during testing

HRV: coefficient of variation based on median

Mean HR significantly decreased at weeks 3 & 4

Mean HRV significantly increased at week 4




Results: Subjective Measures

POMs Vigor significantly decreased at week 4.


Marine anger levels were above normal on POMs.

No changes observed in other subjective measures

Vigor



Anger






Addressing the Baseline Problem

- Scheduling morning sessions for baselines is impossible with USMC at 29 Palms
- Approvals from command were obtained just prior to start of Mojave Viper
- A fully-rested, stress-free baseline may be impossible to obtain from active marines
- Potential solutions
 - Compare to ABM healthy database (n=200) fully-rested & sleep-deprived
 - Acquire data from fully-rested marines when they first arrive at Camp Pendleton



Results: 3C-VT Accuracy & Reaction Time





Conclusions

- Mobile wireless EEG/HR/HRV acquisition technologies provide valid objective measures of fatigue in an operation environment
- Data obtained without interfering in ongoing operations
- EEG/HR/HRV assessed during the 3C-VT quantitatively showed fatigue to be a serious problem in Marines before deployment
- Data indicate that Marines do not self-report fatigue, increasing the need for objective measurements of fatigue



Relevance of Study Results

Fatigue in USMC convoy operations is a serious problem

Estimated 35-50% of US casualties in Iraq occur during convoy attack

Results suggest feasibility of objective fatigue assessment in operational settings

Coupled with appropriate interventions fatigue monitoring could improve safety and save lives



Advanced Brain Monitoring, Inc.

Q & A